

Organizational culture and the adoption of anti-smoking initiatives at small to very small workplaces: An organizational level analysis

Christine M. Kava¹, Edith A. Parker¹, Barbara Baquero¹, Susan J. Curry², Paul A. Gilbert¹, Michael Sauder³, Daniel K. Sewell⁴

ABSTRACT

INTRODUCTION Many workplaces have adopted anti-smoking initiatives to reduce smoking behavior, but small workplaces are less likely to adopt these initiatives. One factor that could influence adoption is organizational culture, defined as the values and assumptions shared by members of an organization. The aim of this study was to examine the types of organizational culture associated with smoking policy strength and adoption of smoking cessation activities at small (20–99 employees) and very small (<20 employees) workplaces. Two study hypotheses were made: An increase in clan culture (characterized by participation in decision-making and human resources development) will be associated with an increase in smoking policy strength (H1) and higher odds of having cessation activities in the workplace (H2).

METHODS Between June and October 2017, executives and employees coming from small and very small workplaces participated in separate surveys. Executives answered questions about their workplace's anti-smoking initiatives, while employees completed a 12-item questionnaire about organizational culture. We aggregated employee data to perform linear and logistic regression at the organizational level.

RESULTS Organizational culture was not significantly associated with smoking policy strength, therefore H1 was not supported. Counter to H2, an increase in clan culture was associated with lower odds of offering smoking cessation activities (OR=0.06; 95% CI: 0.01–0.58).

CONCLUSIONS We did not find support for the hypothesized relationships. External factors and additional cultural characteristics may explain study findings. Continued research on culture and ways to improve tobacco control within smaller workplaces is needed.

AFFILIATION

¹ Department of Community and Behavioral Health, University of Iowa College of Public Health, Iowa, United States

² Department of Health Management and Policy, University of Iowa College of Public Health, Iowa, United States

³ Department of Sociology, University of Iowa, Iowa, United States

⁴ Department of Biostatistics, University of Iowa College of Public Health, Iowa, United States

CORRESPONDENCE TO

Christine M. Kava, Health Promotion Research Center, University of Washington, Box 354804, 1107 NE 45th Street, Suite 400, Seattle, WA 98105, United States. E-mail: ckava@uw.edu

KEYWORDS

smoking, organizational culture, tobacco control, small workplaces

Received: 8 August 2018

Revised: 26 October 2018

Accepted: 29 November 2018

INTRODUCTION

The current smoking rate among US adults is 16%¹. In Iowa, this rate is slightly higher at 17%². Smoking is the leading cause of preventable death, responsible for almost half a million deaths each year³. To combat the negative consequences associated with smoking, workplaces commonly adopt two types of

anti-smoking initiatives: smoke-free policies (rules designed to restrict smoking and protect employees from its harmful effects) and cessation activities (programs designed to reduce smoking). Both have been successful in increasing cessation^{4,5}.

Based on previous studies and definitions^{6–8}, small workplaces are defined here as employing between

20 to 99 employees, and very small workplaces as employing less than 20 employees. These smaller workplaces are less likely to adopt anti-smoking initiatives, with adoption rates increasing as the size of the workplace increases⁷. To reduce disparities in tobacco control and smoking behavior, it is important to understand factors associated with anti-smoking initiative adoption in smaller workplaces.

One factor to consider is organizational culture, defined as the beliefs and assumptions shared by organizational members⁹. Several frameworks exist to conceptualize culture, including the Competing Values Framework (CVF). The CVF has been extensively used in the field of organizational psychology, it is a theoretically driven framework with strong evidence of instrumental reliability and validity^{10,11}, and as a quantitative measure facilitates comparisons of culture across organizations¹².

The CVF posits that there are two dimensions of organizational effectiveness, represented by four culture types: clan, adhocracy, hierarchy, and market¹⁰. Clan culture emphasizes participation in decision-making and human resources development. Adhocracy culture emphasizes innovation and entrepreneurship. Hierarchy culture emphasizes structure and standardization. Market culture emphasizes productivity and competition^{10,13}. Most organizations have a dominant culture type¹⁰, however ‘organizations are expected to reflect all four cultures to some degree’, as noted by Helfrich et al.¹⁴.

Previous studies have found that primary care practices exhibiting a combination of clan and adhocracy cultures were more likely to offer preventative health services, which include services for tobacco¹⁵. In a similar study examining primary care provider adherence to guidelines for treating tobacco use, clan cultures were more likely to adhere to the 5A guidelines (i.e. ask, advise, assess, assist, arrange) for tobacco treatment compared to adhocracy cultures¹⁶. Clan culture is also associated with greater workplace practice satisfaction¹⁷ and well-being of employees¹⁸.

A majority of studies examining the association between culture and health initiative adoption have been conducted in healthcare settings and focused on service delivery. Given the research described above, it could be argued that workplaces that are

higher in clan culture would also be more likely to adopt initiatives for tobacco control. Clan cultures are characterized by a corporate commitment to employees and creation of a ‘humane work environment’¹⁰. These values may extend to health promotion, with leadership taking steps to adopt initiatives that improve the employees’ well-being and facilitate a healthy working environment.

More broadly, culture plays a prominent role in influencing organizational behavior and decision-making¹⁹; as described by Tierney²⁰, ‘an understanding of culture has become essential for those who seek to understand how to foment change in the organization’. Thus, knowing how culture is related to tobacco control can help guide intervention efforts that enhance cultural characteristics associated with initiative adoption. Additional research is also needed among smaller workplaces, which offer unique challenges and opportunities for health promotion²¹. Previous researchers have called for increased efforts to develop health promotion programs feasible for very small workplaces in particular⁶.

The aim of this study was to examine the types of organizational culture associated with smoking policy strength and adoption of smoking cessation activities at small (20–99 employees) and very small (<20 employees) workplaces, in Iowa. We hypothesized the following relationships:

H1: An increase in clan culture will be associated with an increase in smoking policy strength.

H2: An increase in clan culture will be associated with higher odds of having cessation activities in the workplace.

METHODS

Design and recruitment

This study used a cross-sectional survey design. Executives coming from small and very small workplaces in Iowa completed a self-administered online survey about the smoking policies and cessation activities at their workplace. Employees at these same workplaces were then asked to complete a self-administered online survey about their organization’s culture. We identified executives through ReferenceUSA, a business database²². To qualify for inclusion, a business had to be listed as being small or very small, verified as active (open),

and having an executive e-mail address on file. Only one executive per business was contacted. In the event that multiple executives were listed under a business in the database, we contacted the most recent (current) executive.

Based on these criteria, approximately 40000 small and very small businesses were eligible to participate in the study. We initially sampled 2000 businesses from the ReferenceUSA database using quasi-random sampling techniques (e.g. one of every 20th business selected). Based on low response rates, we sampled additional businesses from the database to ensure that our goal of recruiting at least 60 workplaces with executive and employee data was met (we determined this sample size from a power analysis). In total, we sampled approximately 10500 businesses. We stratified businesses by small and very small workplace classification. After removing duplicate records and workplaces previously contacted for a related study (Kava et al.²³), 9470 businesses remained and were subsequently contacted.

We conducted recruitment between June and October 2017. We first sent executives an e-mail about the study, which contained a link to the executive online survey. We contacted non-respondents via e-mail and telephone, up to two times, to again request participation. Executives who completed the survey were sent an e-mail containing a link to the employee survey and a request to distribute this e-mail to their employees. Both groups were eligible to win one of three \$75 gift cards in separate drawings for their participation. We provided all participants with an information sheet containing elements of consent, which included a description of the study's purpose and a statement notifying individuals that participation is voluntary. The University of Iowa's Institutional Review Board approved all study procedures.

Measures

The dependent variables for this study were *smoking policy strength* and *smoking cessation activities offered*. We obtained information on these variables from executives. The independent variable for this study was *organizational culture*, specifically clan culture. Since culture focuses on shared perceptions we asked employees to describe their culture, afterward

aggregating responses to obtain organizational level measures for each culture type. These measures are described in further detail below.

Dependent variables

We calculated a smoking policy strength score (range: 0–9) for each workplace by summing up executive responses to questions on the following: outdoor smoking restrictions (1 item), indoor smoking restrictions (1 item), and additional policies related to tobacco (5 items). The item on outdoor smoking restrictions asked executives to indicate whether employees could smoke outside the building and included three response options ranging from 'Yes, anywhere outside on worksite grounds' (=0) to 'No, nowhere on worksite grounds' (=2). The item on indoor smoking restrictions asked executives to indicate whether employees could smoke inside their workplace building and included three response options ranging from 'Yes, anywhere inside' (=0) to 'No, nowhere inside' (=2). The 2008 Iowa Smokefree Air Act (ISAA) bans smoking in most enclosed areas within places of employment, but there are noted exceptions to this rule (e.g. smoking is allowed on the gaming floor of casinos)²⁴. Thus, we included the item on indoor smoking to ensure that we captured information on all possible workplace smoking policies. Both items were taken from a smoking policy survey developed by the University of Wisconsin's Comprehensive Cancer Center²⁵. We included the five items on additional policies in a checklist and asked executives to indicate whether the policy was present (=1) at their workplace. The items listed went beyond state law requirements, and were included to better assess the effects of culture on policy adoption. An example item is: 'Employees are not allowed to smoke while working, regardless of where they are located (e.g. in the office, working off-site)'.

To calculate smoking cessation activities, executives indicated from a list of items the smoking cessation activities offered by their workplace. An example item is: 'Smoking cessation classes or clinics'. We developed these based on qualitative research findings from a related study²³ and on previous measures^{26–28}. We coded workplaces as either having (=1) or not having (=0) cessation activities.

Independent variable

To assess *organizational culture*, we asked employees to complete a 12-item questionnaire developed by Yeung et al.²⁹, which is based on a validated measure of culture using the CVF¹¹. The measure contains three items corresponding to each of the four culture types, and asks employees to indicate the extent to which each item describes their workplace. Response options range from low (=1) to high (=5). An example of one item corresponding to clan culture: 'This organization is a very personal place. It is like an extended family. People seem to share a lot of themselves'. To create a score for each culture type, corresponding items were averaged together. In the current study, Cronbach's alpha coefficient ranged from 0.73 (hierarchy culture) to 0.79 (clan culture).

Covariates

We adjusted for several workplace characteristics theorized or shown to be associated with tobacco control in previous studies^{7,30,31}. Covariate information was primarily obtained from executives, who reported the percentage of their employees that were employed full-time (0–25%; 26–50%; 51–75%; 76–100%), employees' primary work location (in the building; out in the field), and whether their workplace offered health insurance (yes, no, not sure). We adapted these items from previous studies related to health promotion and smoking cessation^{27,30,32}. Executives also reported their workplace industry based on National Institute for Occupational Safety and Health³³ industry categories.

We used information provided by ReferenceUSA to create two additional variables: workplace size (small; very small) and county urban-rural designation (noncore, micropolitan, small metro, medium metro). County designation categories came from the 2013 National Center for Health Statistics Urban-Rural Classification Scheme for Counties³⁴. We collapsed and recoded variables with smaller cell sizes (≤ 20) based on the distribution of responses: workplace industry (service; non-service), health insurance offered (yes, no/not sure), and percentage of full-time employees (0–75%, >75%).

Data analysis

We conducted data analysis in Stata 15.0³⁵ and R Studio 3.3.2³⁶. We first ran descriptive statistics followed

by bivariate and multivariable analyses. We used linear regression to test H1 and logistic regression to test H2. Given the study's broader aim to assess the relationship between organizational culture and anti-smoking initiative adoption, we included all four culture types in the multivariable regression models. To adjust for potential confounders, we also included the covariates described above. The health insurance variable predicted perfectly cessation activities offered (i.e. all workplaces offering health insurance offered cessation activities), so this covariate was excluded from the analysis.

We aggregated employee data by averaging responses to obtain organizational level measures of culture for each workplace. To justify this aggregation, Rho-within-group coefficients for multi-item scales, $\rho_{wg(j)}$, were calculated for each workplace on the four culture types to assess the extent to which employees at the same workplace agreed on culture³⁷. Mean $\rho_{wg(j)}$ coefficient values were as follows: 0.76 for clan, 0.73 for adhocracy, 0.73 for hierarchy, and 0.78 for market. These values indicate an adequate level of agreement based on previous recommendations³⁸.

RESULTS

Participants

In all, 264 executives, each representing one workplace, and 405 employees participated in this study. Not all workplaces had employee data available. Further, a small number of employees ($n=9$) entered the wrong workplace ID on their survey, which was necessary to link employee responses to those of their executive. In total, executive and employee data were linked at 71 workplaces. Three workplaces employed more than 99 employees, and in two additional cases the executive had recently retired. After excluding executive and employee data from these workplaces, the final sample sizes for this study were 259 executives, 280 employees, and 68 workplaces with both executive and employee data available. The mean number of employees participating at the 68 workplaces was four (range: 1–22).

Main findings

Table 1 provides information on anti-smoking initiatives in the workplace, based on executive reports ($N=259$). Almost all executives reported that their

Table 1. Descriptive statistics: Anti-smoking initiatives (N=259)^a

Variable	Mean	SD
Smoking policy strength score ^b	4.84	1.92
	%	n
Indoor smoking		
Yes, anywhere inside	0.77	2
Yes, but only in designated areas or times	1.93	5
No, nowhere inside	97.30	252
Outdoor smoking		
Yes, anywhere outside	30.62	79
Yes, but only in designated areas or times	31.40	81
No, nowhere outside	37.98	98
Other smoking policies		
Employees not allowed to smoke while working, regardless of location	46.67	119
E-cigarette use is restricted	40.00	102
Smokeless tobacco use is restricted	38.04	97
Workplace has written smoking policy	46.27	118
Other policies	9.80	25
None of the above	19.22	49
One or more activities offered	23.14	59
Cessation activity types		
Referrals for smoking cessation assistance	12.16	31
Incentives for non-smoking or cessation attempts	5.88	15
Self-help materials	8.24	21
Smoking cessation classes or clinics	5.10	13
Lectures or workshops	2.35	6
Other activities	5.88	15
None of the above	76.86	196

a Descriptive statistics calculated using executive data. b Range: 0–9.

workplaces prohibited indoor smoking (97%, n=252). Restrictions on outdoor smoking were less common, with about a third of workplaces (31%, n=79) allowing smoking anywhere outside and 31% (n=81) allowing outdoor smoking in designated areas or times. Approximately 80% of workplaces had additional smoking-related policies. The average smoking policy strength score was 4.84 (SD=1.92) out of nine. Among workplaces offering cessation activities, the most common activity offered was referral for smoking cessation assistance (12%, n=31).

Table 2 describes the organizational characteristics of the sample, with employees at 68 workplaces providing data on culture. Mean culture scores ranged from 3.12 (SD=0.69) for adhocracy

Table 2. Descriptive statistics: Other organizational characteristics (N=259)^a

Variable	Mean	SD ^b
Culture		
Clan	3.81	0.68
Adhocracy	3.12	0.69
Hierarchical	3.37	0.62
Market	3.72	0.51
	%	n
Workplace industry		
Service	53.91	138
Non-service	46.09	118
Workplace size		
Very small (<20 employees)	57.92	150
Small (20–99 employees)	42.08	109
Employee work location		
In the building	81.71	210
In the field	18.29	47
Health insurance offered		
Yes	69.26	178
No/not sure	30.74	79
Insurance coverage for cessation medication^c		
Yes	27.98	47
No	17.86	30
Not sure	54.17	91
Per cent of full-time employees		
0–75%	42.97	110
>75%	57.03	146
County area designation		
Non-core	31.35	79
Micropolitan	13.10	33
Small metro	21.83	55
Medium metro	33.73	85

a Culture calculated only among workplaces with employee data available (n=68).

b SD: standard deviation. c Asked only among employers whose workplace offered health insurance (n=178).

culture to 3.81 (SD=0.68) for clan culture. In bivariate analysis (data not shown in table), correlations among the four culture types ranged from 0.136 to 0.645. Culture was not significantly associated with smoking policy strength. Compared to small workplaces, very small workplaces had a lower mean smoking policy strength score (p=0.012). The mean score for clan culture was lower among workplaces offering cessation activities (p=0.000).

Table 3. Linear regression on smoking policy strength (n=63)

Variable	β	SE ^a	95% CI ^b	p
Culture				
Clan	-0.13	0.53	(-1.21, 0.94)	0.806
Adhocracy	-0.42	0.55	(-1.53, 0.69)	0.448
Hierarchical	0.37	0.48	(-0.60, 1.34)	0.444
Market	-0.20	0.64	(-1.49, 1.10)	0.761
Workplace industry (ref: Non-service)				
Service	-0.37	0.58	(-1.54, 0.80)	0.530
Workplace size (ref: Very small)				
Small	0.84	0.57	(-0.31, 1.99)	0.148
Work location (ref: In the field)				
In the building	0.85	0.70	(-2.25, 0.56)	0.232
Health insurance offered (ref: Yes)				
No/not sure	0.97	0.69	(-0.42, 2.37)	0.168
Per cent full-time employees (ref: 0–75%)				
>75%	0.75	0.62	(-0.48, 1.99)	0.228
County designation (ref: Non-core)				
Micropolitan	0.00	0.74	(-1.49, 1.48)	0.997
Small metro	-0.67	0.80	(-2.27, 0.94)	0.408
Medium metro	0.02	0.67	(-1.32, 1.37)	0.972
Constant	6.29	2.35	(1.57, 11.01)	0.010*
R²	0.18			

a SE: standard error. b CI: confidence interval. *p<0.05, **p<0.01, ***p<0.001.

Table 3 shows the results for the multivariable linear regression predicting smoking policy strength. No significant relationship existed between clan culture and smoking policy strength (p=0.806), therefore H1 was not supported. No other culture types were significantly associated with smoking policy strength. After adjusting for workplace characteristics, differences in smoking policy strength by workplace size were no longer significant. A *post hoc* analysis excluding the additional policies in the calculation of the smoking policy strength variable revealed the same findings (data not shown in table). Table 4 shows the findings from the logistic regression predicting smoking cessation activities. As clan culture increased, the odds of having smoking cessation activities decreased (OR=0.06; 95% CI: 0.01–0.58; p=0.015). These findings do not support H2, which theorized that an increase in clan culture would be associated with *higher* odds of offering cessation activities. No other culture types were associated with the likelihood of offering cessation activities.

Table 4. Logistic regression on smoking cessation activities (n=63)

Variable	OR ^a	95% CI ^b	p
Culture			
Clan	0.06	(0.01, 0.58)	0.015*
Adhocracy	1.09	(0.13, 9.39)	0.937
Hierarchical	2.71	(0.47, 15.56)	0.263
Market	1.72	(0.22, 13.24)	0.600
Workplace industry (ref: Non-service)			
Service	0.48	(0.09, 2.64)	0.398
Workplace size (ref: Very small)			
Small	0.68	(0.12, 3.83)	0.659
Work location (ref: In the field)			
In the building	0.09	(0.01, 0.79)	0.030*
Per cent full-time employees (ref: 0–75%)			
>75%	0.61	(0.10, 3.87)	0.598
County designation (ref: Non-core)			
Micropolitan	3.85	(0.36, 40.76)	0.263
Small metro	0.62	(0.04, 9.60)	0.731
Medium metro	2.72	(0.31, 24.08)	0.369

a OR: odds ratio. b CI: confidence interval. *p<0.05, **p<0.01, ***p<0.001.

DISCUSSION

While previous studies have examined organizational culture's relationship to healthcare outcomes, this study contributes new knowledge on the association between culture and health promotion initiatives related to tobacco. Applying a framework commonly used in organizational psychology (CVF) to gain a greater understanding of tobacco control at smaller workplaces, this study also addressed these relationships within the context of a statewide smoking ban. This study found no significant relationship between clan culture and smoking policy strength. Contrary to expectations, workplaces stronger in clan culture were less likely to offer cessation activities. This is inconsistent with previous studies that have found positive relationships between clan culture and health practices^{15,16}.

The lack of support for our study's hypotheses may be due to several reasons. This study attempted to assess policy strength within the context of the ISAA, a law prohibiting smoking at most workplaces in Iowa²⁴. By including questions on initiatives not covered by this Act and creating a total score based on all policy questions, we attempted to gain a better understanding of how culture may influence policy adoption independently. However, given the ISAA's longstanding presence (10 years) and declining acceptability of smoking in public places, it could be that culture has less of an influence on a workplace's decision to adopt anti-smoking initiatives. In a related qualitative study assessing the barriers and facilitators to anti-smoking initiative adoption among smaller workplaces²³, several participants described the ISAA as a reason for policy adoption. Some participants also described wanting to create a healthy work environment, which may indicate that there are additional aspects of organizational culture (e.g. health culture) more important to policy adoption than culture more broadly.

Regarding the significant and negative association between clan culture and cessation activities offered, managers may not perceive a lack of need for activities, particularly if smoking employee rates are low. This idea is supported by qualitative data from a related study, where several participants described this lack of need²³. At the same time, managers may be willing to offer activities if a perceived need from employees arose. This may be

especially true for workplaces strong in clan culture, characterized by employee participation in decision-making processes. Similarly, the supportive and interpersonal nature of clan culture organizations could make managers less willing to provide advice to employees about their health. This unwillingness may stem from manager fears of being viewed as paternalistic, a notion that contrasts with the characteristics of clan culture and a cited barrier to health promotion in previous studies³⁹.

Bivariate analysis revealed a lower smoking policy strength score among very small workplaces compared to small workplaces. This relationship became non-significant after accounting for other organizational characteristics, suggesting that differences in these characteristics are important to policy adoption and warrant further attention. Differences by workplace size in smoking cessation activity adoption did not exist, although overall the number of workplaces offering these activities was low (23%, $n=59$). Given this information, continued efforts to develop tobacco control programs tailored to the environment of smaller workplaces are needed.

Several potential limitations warrant consideration. A lack of generalizability and power to detect small effect sizes may be present, as the response rate and sample sizes for this study were low. While previous studies have used ReferenceUSA and similar databases for recruitment⁴⁰⁻⁴², others have called into question the validity of these databases⁴³. In the current study, approximately 20% of the recruitment e-mails that were sent to executives came back as undeliverable, suggesting a lack of accuracy that could help to explain the low response rates we observed. Since all workplaces included in this study came from Iowa, they may not be representative of all workplaces in the US. In states with less comprehensive legislation or different social norms related to smoking, these relationships may differ. Nevertheless, our findings are likely to be relevant to other workplaces in similar settings and in states with comparable legislation on tobacco control, and contribute to our limited understanding of these relationships within smaller workplaces.

Executives and employees who did not perceive smoking as an issue at their workplace may have

been less likely to participate in the study, leading to non-response bias. After receiving an invitation to participate in the study, some executives indicated that they did not feel their workplace qualified to participate, either because their workplace was smoke-free or because they did not currently employ any smokers. To reduce the potential for bias, these individuals were informed by the first author (CMK) that we were seeking to recruit a diverse range of workplaces. We also adapted our recruitment materials to reflect this point. Lastly, while organizational culture focuses on *shared* assumptions⁴⁴, some of the workplaces included in this study had only one employee participant. Given our focus on smaller workplaces this was not unexpected, but makes it harder to understand culture from a shared viewpoint.

Previous studies have described the challenges associated with measuring culture, a complex construct with multiple dimensions. As described by Scott et al.⁴⁵, 'A problem with trying to assess highly complex phenomena like culture is that experts rarely agree on which are the essential dimensions to measure'. They go on to suggest assessments of culture via qualitative or mixed methods approaches, which may be better at capturing culture's complexity. For example, it was speculated earlier that the supportive and interpersonal nature of organizations strong in clan culture may make these workplaces less likely to offer activities for cessation. Taking a qualitative research approach would allow a more detailed exploration of the nuances of clan culture that influence decision-making, leading to a better understanding of how to increase tobacco control within smaller workplaces to improve employee health.

CONCLUSIONS

The current study found no association between organizational culture and smoking policy strength. In a finding counter to H2, workplaces stronger in clan culture were less likely to offer activities to help their employees quit smoking. State-wide smoking bans and additional cultural characteristics may help to explain organizational decision-making related to tobacco control. After accounting for several organizational features, no differences in anti-smoking initiative adoption by workplace size existed. Adoption of

cessation activities was low overall, suggesting a need for increased tobacco control in smaller workplaces.

REFERENCES

1. Jamal A, Phillips E, Gentzke AS, et al. Current cigarette smoking among adults - United States, 2016. *Morb Mortal Weekly Rep.* 2018;67(2):53-59. doi:10.15585/mmwr.mm6702a1
2. Centers for Disease Control and Prevention. Map of current cigarette use among adults 2016. <https://www.cdc.gov/statesystem/cigaretteuseadult.html>. Accessed November 16, 2016.
3. U.S. Department of Health and Human Services. The health consequences of smoking -- 50 years of progress: A report of the Surgeon General. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2014.
4. Bauer JE, Hyland A, Li Q, Steger C, Cummings KM. A longitudinal assessment of the impact of smoke-free worksite policies on tobacco use. *Am J Public Health.* 2005;95(6):1024-1029. doi:10.2105/AJPH.2004.048678
5. Cahill K, Moher M, Lancaster T. Workplace interventions for smoking cessation. *Cochrane Database of Systematic Reviews.* 2008(4):CD003440. doi:10.1002/14651858.CD003440.pub4
6. Harris JR, Hannon PA, Beresford SA, Linnan LA, McLellan DL. Health promotion in smaller workplaces in the United States. *Annu Rev Public Health.* 2014;35:327-342. doi:10.1146/annurev-publhealth-032013-182416
7. Linnan L, Bowling M, Childress J, et al. Results of the 2004 National Worksite Health Promotion Survey. *Am J Public Health.* 2008;98(8):1503-1509. doi:10.2105/AJPH.2006.100313
8. Headd B. The characteristics of small-business employees. *Mon Labor Rev.* 2000;123(4):13-18.
9. Butterfoss FD, Kegler MC, Francisco VT. Mobilizing organizations for health promotion: Theories of organizational change. In: Glanz K, Rimer BK, Viswanath K, eds. *Health behavior and health education: Theory, research, and practice.* San Francisco, CA: Jossey-Bass; 2008:335-357.
10. Cameron KS, Quinn RE. Diagnosing and changing organizational culture: Based on the Competing Values Framework. San Francisco, CA: Jossey-Bass; 2006.
11. Quinn RE, Spreitzer GM. The psychometrics of the competing values culture instrument and an analysis of the impact of organizational culture on quality of life. *Research in Organizational Change and Development.* 1991;5:115-142.
12. Yauch CA, Steudel HJ. Complementary use of qualitative and quantitative cultural assessment methods. *Organ Res Methods.* 2003;6(4):465-481. doi:10.1177/1094428103257362
13. Quinn RE, Rohrbaugh J. A spatial model of effectiveness

- criteria: Towards a Competing Values approach to organizational analysis. *Manag Sci.* 1983;29(3):363-377. doi:10.1287/mnsc.29.3.363
14. Helfrich CD, Li YF, Mohr DC, Meterko M, Sales AE. Assessing an organizational culture instrument based on the Competing Values Framework: exploratory and confirmatory factor analyses. *Implement Sci.* 2007;2:13. doi:10.1186/1748-5908-2-13
 15. Hung DY, Rundall TG, Tallia AF, Cohen DJ, Halpin HA, Crabtree BF. Rethinking prevention in primary care: Applying the chronic care model to address health risk behaviors. *Milbank Q.* 2007;85(1):69-91. doi:10.1111/j.1468-0009.2007.00477.x
 16. Hung DY, Leidig R, Shelley DR. What's in a setting?: Influence of organizational culture on provider adherence to clinical guidelines for treating tobacco use. *Health Care Manage Rev.* 2014;39(2):154-163. doi:10.1097/HMR.0b013e3182914d11
 17. Zazzali JL, Alexander JA, Shortell SM, Burns LR. Organizational culture and physician satisfaction with dimensions of group practice. *Health Serv Res.* 2007;42(3 Pt 1):1150-1176. doi:10.1111/j.1475-6773.2006.00648.x
 18. Marchand A, Haines VY 3ed, Dextras-Gauthier J. Quantitative analysis of organizational culture in occupational health research: A theory-based validation in 30 workplaces of the organizational culture profile instrument. *BMC Public Health.* 2013;13:443. doi:10.1186/1471-2458-13-443
 19. Schein EH. Organizational culture. *Am Psychol* 1990;45(2):109-119. doi:10.1037/0003-066x.45.2.109
 20. Tierney WG. The impact of culture on organizational decision making: Theory and practice in higher education. 1st ed. Sterling, VA: Stylus Publishing, LLC; 2008.
 21. McCoy K, Stinson K, Scott K, Tenney L, Newman LS. Health promotion in small business: A systematic review of factors influencing adoption and effectiveness of worksite wellness programs. *J Occup Environ. Med* 2014;56(6):579-587. doi:10.1097/JOM.0000000000000171
 22. ReferenceUSA. Home page. ReferenceUSA. <http://www.referenceusa.com/Home/Home>. Accessed August 9, 2016.
 23. Kava CM, Parker EA, Baquero B, et al. A qualitative assessment of the smoking policies and cessation activities at smaller workplaces. *BMC Public Health* 2018;18(1):1094. doi:10.1186/s12889-018-6001-9
 24. The Iowa Legislature. CHAPTER 142 D SMOKEFREE AIR ACT. <https://coolice.legis.iowa.gov/Cool-ICE/default.asp?category=billinfo&service=IowaCode&ga=83&input=142D>. Accessed February 21, 2017.
 25. Anderson AE, Moberg DP, Hill B, Christiansen A. 2001 Wisconsin Worksite Smoking Policy Survey. 2002. Accessed May 30, 2017.
 26. Centers for Disease Control and Prevention. Workplace health site visit interview questions. <http://www.cdc.gov/workplacehealthpromotion/pdf/workplacehealthinterviewquestions.pdf>. Accessed 30 Sep, 2016.
 27. Centers for Disease Control and Prevention. Worksite ScoreCard. <https://www.cdc.gov/workplacehealthpromotion/initiatives/healthscorecard/worksite-scorecard.html>. Accessed May 30, 2017.
 28. Flynn BS, Gurdon MA, Secker-Walker RH. Cigarette smoking control strategies of firms with small work forces in two Northeastern States. *Am J Health Promot.* 1995;9(3):202-209, 219. doi:10.4278/0890-1171-9.3.202
 29. Yeung AK, Brockbank JW, Ulrich D. Organizational culture and human resource practices: An empirical assessment. *Research in Organizational Change and Development.* 1991;5:59-81.
 30. Gottlieb NH, Lovato CY, Weinstein R, Green LW, Eriksen MP. The implementation of a restrictive worksite smoking policy in a large decentralized organization. *Health Educ Q.* 1992;19(1):77-100. doi:10.1177/109019819201900106
 31. Hahn EJ, Rayens MK, York N. Readiness for smoke-free policy and overall strength of tobacco control in rural tobacco-growing communities. *Health Promot Pract.* 2013;14(2):238-246. doi:10.1177/1524839912446479
 32. Eby LT, Laschober TC, Muilenburg JL. Understanding counselors' implementation of tobacco cessation services with patients. *J Subst Abuse Treat.* 2014;47(5):314-320. doi:10.1016/j.jsat.2014.06.007
 33. National Institute for Occupational Safety and Health. Sectors. <https://www.cdc.gov/niosh/nora/sectorapproach.html>. Accessed October 1, 2017.
 34. Ingram D, Franco S. 2013 NCHS urban-rural classification scheme for counties. National Center for Health Statistics. *Vital Health Stat.* 2014;2(166).
 35. Stata Statistical Software: Release 15 [computer program]. College Station, TX: StataCorp LLC; 2017.
 36. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. The R Foundation; 2017. <http://www.R-project.org/>. Accessed September 29, 2017.
 37. James LR, Demaree RG, Wolf G. Estimating within-group interrater reliability with and without response bias. *J Appl Psychol.* 1984;69(1):85-98. doi:10.1037/0021-9010.69.1.85
 38. Klein KJ, Kozlowski SWJ. From micro to meso: Critical steps in conceptualizing and conducting multilevel research. *Organ Res Methods.* 2000;3(3):211-236. doi:10.1177/109442810033001
 39. Eakin JM. Leaving it up to the workers: Sociological perspective on the management of health and safety in small workplaces. *Int J Health Serv.* 1992;22(4):689-704. doi:10.2190/dnv0-57vv-fj7k-8ku5
 40. Usan K, Forster JL, Chen V. Smoking policies in small worksites in Minnesota. *Am J Health Promot.* 2007;21(5):416-421. doi:10.4278/0890-1171-21.5.416

41. Hannon PA, Harris JR, Sopher CJ, et al. Improving low-wage, mid-sized employers' health promotion practices: A randomized controlled trial. *Am J Prev Med*. 2012;43(2):125-133. doi:10.1016/j.amepre.2012.04.014
42. DeJoy DM, Dyal MA, Padilla HM, Wilson MG. National workplace health promotion surveys: The Affordable Care Act and future surveys. *Am J Health Promot*. 2014;28(3):142-145. doi: 10.4278/ajhp.121212-CIT-602
43. Cook RG, Campbell DK, Kelly C. An issue of trust: Are commercial databases really reliable? *Journal of Business & Finance Librarianship* 2012;17(4):300-312. doi:10.1080/08963568.2012.712501
44. Schein EH. *Organizational culture and leadership: A dynamic view*. San Francisco, CA: Jossey-Bass; 1985.
45. Scott T, Mannion R, Davies H, Marshall M. The quantitative measurement of organizational culture in health care: A review of the available instruments. *Health Serv Res*. 2003;38(3):923-945. doi:10.1111/1475-6773.00154

CONFLICTS OF INTEREST

The authors declare that they have no competing interests, financial or otherwise, related to the current work. D. K. Sewell reports grants from US Department of Health & Human Services, Centers for Disease Control & Prevention, during the conduct of the study. The rest of the authors also have completed and submitted an ICMJE form for disclosure of potential conflicts of interest.

FUNDING

This work was supported by Cooperative Agreement Number 1 U48 DP005021-01 from the Centers for Disease Control and Prevention. The findings and conclusions in this article are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention. This work was also supported by funding from the University of Iowa Graduate & Professional Student Government student research grant, and University of Iowa College of Public Health Advancing Graduate Student Success Award. The funding bodies were not involved in the study design, collection, analysis, and interpretation of data, or in writing the manuscript.

PROVENANCE AND PEER REVIEW

Not commissioned; externally peer reviewed.